

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

MXI1 monoclonal antibody (M08), clone 1F3

Catalog Number: H00004601-M08

Regulation Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody

raised against a partial recombinant MXI1.

Clone Name: 1F3

Immunogen: MXI1 (NP $_$ 001008541.1, 73 a.a. \sim 182 a.a) partial recombinant protein with GST tag. MW of the

GST tag alone is 26 KDa.

Sequence:

LEEAERKSQHQLENLEREQRFLKWRLEQLQGPQEME RIRMDSIGSTISSDRSDSEREEIEVDVESTEFSHGEVD NISTTSISDIDDHSSLPSIGSDEGYSSASVKLSFTS

Host: Mouse

Reactivity: Human

Applications: ELISA, WB-Re, WB-Tr

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Isotype: IgG2a Kappa

Storage Buffer: In 1x PBS, pH 7.4

Storage Instruction: Store at -20°C or lower. Aliquot to

avoid repeated freezing and thawing.

Entrez GenelD: 4601

Gene Symbol: MXI1

Gene Alias: MAD2, MGC43220, MXD2, MXI, bHLHc11

Gene Summary: Expression of the c-myc gene, which produces an oncogenic transcription factor, is tightly regulated in normal cells but is frequently deregulated in human cancers. The protein encoded by this gene is a

transcriptional repressor thought to negatively regulate MYC function, and is therefore a potential tumor suppressor. This protein inhibits the transcriptional activity of MYC by competing for MAX, another basic helix-loop-helix protein that binds to MYC and is required for its function. Defects in this gene are frequently found in patients with prostate tumors. Three alternatively spliced transcripts encoding different isoforms have been described. Additional alternatively spliced transcripts may exist but the products of these transcripts have not been verified experimentally. [provided by RefSeq]